

Claremont Colleges Scholarship @ Claremont

All HMC Faculty Publications and Research

HMC Faculty Scholarship

2-1-1994

Uniqueness of Stable and Unstable Positive Solutions for Semipositone Problems

Alfonso Castro
Harvey Mudd College

Sudhasree Gadam
University of North Texas

Recommended Citation

Castro, Alfonso and Sudhasree Gadam. "Uniqueness of the stable and unstable positive solutions for semipositone problems", *Nonlinear Analysis T.M.A.*, Vol. 22, No. 4 (1994), pp. 425-429.

This Article - postprint is brought to you for free and open access by the HMC Faculty Scholarship at Scholarship @ Claremont. It has been accepted for inclusion in All HMC Faculty Publications and Research by an authorized administrator of Scholarship @ Claremont. For more information, please contact scholarship@cuc.claremont.edu.



UNIQUENESS OF STABLE AND UNSTABLE POSITIVE SOLUTIONS FOR SEMIPOSITONE PROBLEMS

ALFONSO CASTRO and GADAM SUDHASREE

Department of Mathematics, University of North Texas, Denton, TX 76203-5116, U.S.A.

(Received 20 October 1992; received for publication 2 June 1993)

Key words and phrases: Semipositone problems, stable and unstable positive solutions.

1. INTRODUCTION

IN THIS PAPER we consider positive solutions to the equation

$$-\Delta u(x) = \lambda f(u(x)) \quad \text{for } x \in \Omega \quad (1.1)$$

$$u(x) = 0 \quad \text{for } x \in \partial\Omega \quad (1.2)$$

where Ω denotes the unit ball in \mathbb{R}^N ($N > 1$), centered at the origin and $\lambda > 0$. Here $f: [0, \infty) \rightarrow \mathbb{R}$ is assumed to be monotonically increasing, concave and such that

$$f(0) < 0 \text{ (semipositone)}, \quad f(u) > 0 \text{ for some } u > 0. \quad (1.3)$$

Let F be defined by $F(t) = \int_0^t f(s) ds$. We let β and θ denote the unique positive zeros of f and F , respectively. It can be easily seen that if u is a positive solution to (1.1), (1.2) then $u(0) > \theta$. We prove that for each λ there is at most one stable and one unstable positive solution to the problem (1.1), (1.2). More precisely, we prove the following theorem.

THEOREM A. If f is as above and $\lim_{t \rightarrow \infty} f'(t) = 0$, then there exist λ_1 and λ_2 with $\lambda_1 > \lambda_2$ such that for $\lambda > \lambda_1$ the problem (1.1), (1.2) has a unique positive solution, which is stable; for $\lambda \in (\lambda_2, \lambda_1]$ it has exactly two positive solutions, one stable and one unstable; and for $\lambda = \lambda_2$ it has a unique positive solution.

We note that the hypothesis $\lim_{t \rightarrow \infty} f'(t) = 0$ is also necessary for the existence of positive solutions for large values of λ (see [1, lemma 2.3]). The case $N = 1$ can be found in [2]. For other results in the radial case the reader is referred to [1, 3, 4]. In contrast with the positone case ($f(0) > 0$), theorem A shows the nonuniqueness of positive solutions for the semipositone case. In a forthcoming paper we prove that for any $\lambda > 0$, the problem (1.1), (1.2) has at most one positive solution when f is convex.

The positive solutions to (1.1), (1.2) are known to be radially symmetric (see [5]). Thus, the problem reduces to the study of solutions to the ordinary differential equation

$$u'' + ((N-1)/r)u' + \lambda f(u) = 0 \text{ in } (0, 1), \quad (1.4)$$

$$u'(0) = 0 \quad \text{and} \quad u(1) = 0, \quad (1.5)$$

where $'$ denotes differentiation with respect to $r = \|x\|$. Our proofs use the one-dimensional maximum principle, eigenvalue comparison arguments, and the bifurcation analysis at a degenerate solution.

2. FIRST AND SECOND VARIATIONS WITH RESPECT TO PARAMETERS

For any $d > 0$, let $u(r, \lambda, d)$ denote the solution to the initial value problem

$$u'' + ((N-1)/r)u' + \lambda f(u) = 0 \quad (2.1)$$

$$u'(0) = 0 \quad \text{and} \quad u(0) = d. \quad (2.2)$$

Thus, $u(r, \lambda, d)$ is a positive solution to (1.1), (1.2) if, in addition, it satisfies

$$u(r, \lambda, d) > 0 \quad \text{for } r \in (0, 1) \text{ and } u(1, \lambda, d) = 0. \quad (2.3)$$

Let v denote the solution to the corresponding linearized problem

$$v'' + ((N-1)/r)v' + \lambda f'(u)v = 0 \quad (2.4)$$

$$v(0) = 1 \quad \text{and} \quad v'(0) = 0. \quad (2.5)$$

Let w denote the solution to the problem

$$w'' + ((N-1)/r)w' + \lambda f'(u)w = -\lambda f''(u)v^2 \quad (2.6)$$

$$w(0) = 0 \quad \text{and} \quad w'(0) = 0. \quad (2.7)$$

That is, v is the derivative of $u(r, \lambda, d)$ with respect to d and w is the second derivative of $u(r, \lambda, d)$ with respect to d .

LEMMA 1. If u is a solution to (2.1)–(2.3) then v has at most one zero in $[0, 1]$.

Proof. Let u be a solution to (2.1)–(2.3) such that v changes sign in $[0, 1]$. Let s be the first zero of v . That is $v > 0$ in $[0, s)$ and $v(s) = 0$. Let $r_0 \in (0, 1)$ be such that $u(r_0) = \beta$. Such an r_0 exists since $u(0) > \theta$ and $u(1) = 0$. We first prove that $s \notin (0, r_0)$. Suppose, on the contrary, that $s \in (0, r_0)$. By setting

$$\varphi(r) = v(r)/f(u(r)),$$

we obtain that in $(0, s)$, φ satisfies

$$\varphi'' + (((N-1)/r) + (2f'(u)u'/f(u)))\varphi' + (f''(u)(u')^2/f(u))\varphi = 0, \quad (2.8)$$

$$\varphi'(0) = 0, \quad \varphi(0) > 0, \quad \varphi(s) = 0. \quad (2.9)$$

Since $f(u(r)) > 0$ in $(0, s)$, and $f'' \leq 0$, by the maximum principle (see [6, theorem 4, p. 7]) we conclude that φ attains its maximum at 0 and $\varphi'(0) < 0$. Since this contradicts (2.9) we see that $s \geq r_0$.

Now we rule out the possibility of v having a second zero in $[r_0, 1]$. Suppose v has two zeros in $[r_0, 1]$. Let s_1 and s_2 be the first two zeros of v . Let $t \in (s_1, s_2)$ be such that $v'(t) = 0$ and $v' > 0$ in (t, s_2) . Multiplying (2.4) by u' and integrating over (t, s_2) we get

$$u'(s_2)v'(s_2) - \int_t^{s_2} v'u'' + \int_t^{s_2} ((N-1)/r)v'u' + \lambda \int_t^{s_2} f'(u)u'v = 0.$$

This, with (2.1), yields

$$u'(s_2)v'(s_2) + 2 \int_t^{s_2} ((N-1)/r)v'u' + \lambda \int_t^{s_2} (f(u)v)' = 0.$$

Hence, we get

$$u'(s_2)v'(s_2) - \lambda f(u(t))v(t) > 0,$$

which is a contradiction to our assumption that $t \in (r_0, 1)$. Thus v cannot have a second zero in $(0, 1]$. This proves the lemma. ■

LEMMA 2. If $u(\cdot, \lambda_0, d_0)$ is a solution to (2.1)–(2.3) and $v(1, \lambda_0, d_0) = 0$, then $u_\lambda(1, \lambda_0, d_0) < 0$ and $u_{dd}(1, \lambda_0, d_0) > 0$. Moreover, there exists an $\varepsilon > 0$ and a differentiable function $\Lambda: (d_0 - \varepsilon, d_0 + \varepsilon) \rightarrow \mathbb{R}$ such that for any $d \in (d_0 - \varepsilon, d_0 + \varepsilon)$, $u(\cdot, \Lambda(d), d)$ is a solution to (2.1)–(2.3), $\Lambda'(d_0) = 0$ and $\Lambda''(d_0) > 0$. In addition, if $u(1, \lambda, d) = 0$ with $|d - d_0| < \varepsilon$, $|\lambda - \lambda_0| < \varepsilon$ then $\lambda = \Lambda(d)$. In particular, if $|\lambda - \lambda^*| < \varepsilon$, $|d - d^*| < \varepsilon$ then $\lambda \in (\lambda_0, \lambda_0 + \varepsilon)$.

Proof. An elementary rescaling gives that

$$u(r/\rho, \lambda, d) = u(r, \lambda/\rho^2, d)$$

for all $\rho > 0$. Differentiating this with respect to ρ and evaluating at $\rho = 1$ we obtain

$$ru'(r, \lambda, d) = 2\lambda u_\lambda(r, \lambda, d). \quad (2.10)$$

By lemma 1, $v > 0$ on $[0, 1)$. Thus, v is an eigenfunction corresponding to the smallest eigenvalue of the problem

$$\begin{aligned} -\Delta\varphi - \lambda f'(u)\varphi &= \mu_1\varphi & \text{in } \Omega, \\ \varphi &= 0 & \text{on } \partial\Omega. \end{aligned}$$

If $u'(1, \lambda, d_0) = 0$, then $\partial u / \partial x_i$, $1 \leq i \leq N$ are also eigenfunctions corresponding to the eigenvalue $\mu_1 = 0$. Since this contradicts the fact that μ_1 is simple we have $u'(1, \lambda_0, d_0) < 0$. This and (2.10) give $u_\lambda(1, \lambda_0, d_0) < 0$.

By the implicit function theorem there exists an $\varepsilon > 0$ and a differentiable function $\Lambda: (d_0 - \varepsilon, d_0 + \varepsilon) \rightarrow \mathbb{R}$ such that for any $d \in (d_0 - \varepsilon, d_0 + \varepsilon)$, $u(1, \Lambda(d), d) = 0$. In particular $u(\cdot, \Lambda(d), d)$ satisfies (2.1)–(2.3). Differentiating $u(1, \Lambda(d), d) = 0$ with respect to d , we have

$$u_\lambda(1, \Lambda(d_0), d_0)\Lambda'(d_0) + u_d(1, \Lambda(d_0), d_0) = 0. \quad (2.11)$$

This, (2.10) and the assumption $v(1, \lambda_0, d_0) = 0$ imply that $\Lambda'(d_0) = 0$. Differentiating again with respect to d , we obtain

$$u_\lambda(1, \Lambda(d_0), d_0)\Lambda''(d_0) + u_{dd}(1, \Lambda(d_0), d_0) = 0, \quad (2.12)$$

where we have used that $\Lambda'(d_0) = 0$. Multiplying (2.4) by $r^{N-1}w$ and (2.6) by $r^{N-1}v$, subtracting one from the other and integrating by parts on $[0, t)$ we see that

$$t^{n-1}(v'(t)w(t) - w'(t)v(t)) = \lambda \int_0^t r^{N-1}f''(u(r))v^3(r) dr.$$

Now since by lemma 1 $u_d(r, \lambda_0, d_0) > 0$ on $[0, 1)$, it follows that $u_{dd}(1, \lambda_0, d_0) > 0$. This, in turn, implies that $\Lambda''(d_0) > 0$ which proves the lemma. ■

LEMMA 3. If $\Gamma \subset \mathbb{R}^2$ is a component of $\{(\lambda, d): u(\cdot, \lambda, d) \text{ is a solution to (2.1)–(2.3)}\}$, then Γ is unbounded in the λ direction.

Proof. Let $(\lambda_0, d_0) \in \Gamma$. We distinguish three cases, namely $u_d(1, \lambda_0, d_0) > 0$, $u_d(1, \lambda_0, d_0) = 0$ and $u_d(1, \lambda_0, d_0) < 0$.

If $u_d(1, \lambda_0, d_0) > 0$, by the implicit function theorem there exists a $\delta > 0$ and an increasing differentiable function $\zeta: (\lambda_0 - \delta, \lambda_0 + \delta) \rightarrow \mathbb{R}$ such that

$$u(1, \lambda, \zeta(\lambda)) = 0. \quad (2.13)$$

Hence, $u(\cdot, \lambda, \zeta(\lambda))$ satisfies (2.1)–(2.3). We claim that ζ defined by (2.13) can be extended to (λ, ∞) . In fact, suppose $\lambda^* = \sup\{\lambda: \zeta \text{ can be extended to } (\lambda_0, \lambda) \text{ with } u(\cdot, s, \zeta(s)) \text{ satisfying (2.1)–(2.3)}\} < \infty$. Letting $d^* := \sup\{\zeta(\lambda): \lambda < \lambda^*\}$, we see that $u_d(1, \lambda^*, d^*) = 0$. Taking ε as in lemma 2 we have a contradiction because if $|\lambda - \lambda^*| < \varepsilon$, $|d - d^*| < \varepsilon$ are such that $u(\cdot, \lambda, d)$ satisfies (2.1)–(2.3) then $\lambda > \lambda^*$. Thus, $\lambda^* = \infty$, which proves that Γ is unbounded in the λ direction.

If $u_d(1, \lambda_0, d_0) = 0$, applying lemma 2 we see that $u(\cdot, \Lambda(d_0 + \varepsilon/2), d_0 + \varepsilon/2)$ is in Γ and $u_d(1, \Lambda(d_0 + \varepsilon/2), d_0 + \varepsilon/2) > 0$. Thus, by the arguments in the latter paragraph, we conclude that Γ is unbounded in the λ direction.

Finally, if $u_d(1, \lambda_0, d_0) < 0$, then by the implicit function theorem there exists a $\delta > 0$ and a decreasing differentiable function $\eta: (\lambda_0 - \delta, \lambda_0 + \delta) \rightarrow \mathbb{R}$ such that $u(1, \lambda, \eta(\lambda)) = 0$ and $u(\cdot, \lambda, \eta(\lambda))$ satisfies (2.1)–(2.3). Let $\lambda^* = \inf\{\lambda: \eta \text{ can be extended to } (\lambda, \lambda_0) \text{ with } u(\cdot, s, \eta(s)) \text{ satisfying (2.1)–(2.3)}\}$. Letting $d^* = \sup\{\eta(\lambda): \lambda < \lambda_0\}$ we see that $u(\cdot, \lambda^*, d^*)$ satisfies (2.1)–(2.3). Because $f(0) < 0$ for $\lambda > 0$ near zero the problem (2.1)–(2.3) does not have a solution. In particular, $\lambda^* > 0$ and $d^* < \infty$. Since $u_d(1, \lambda^*, d^*) = 0$, arguing as in the previous case, we see that Γ is unbounded in the λ direction which proves the lemma. ■

3. PROOF OF THEOREM A

First we show that $\Gamma = \{(\lambda, d): u(\cdot, \lambda, d) \text{ satisfies (2.1)–(2.3)}\}$ is connected. In fact, if Γ_1 and Γ_2 are two connected components then by lemma 3 both contain elements of the form (λ, d) with $\lambda > 0$ large. However, by theorem A of [1] for λ large (2.1)–(2.3) has a unique solution. Since this contradicts that Γ_1 and Γ_2 are disjoint, we have $\Gamma_1 = \Gamma_2 = \Gamma$.




Fig. 1.

Next, we show that there exists a unique $(\lambda_2, d_2) \in \Gamma$ such that $u_d(1, \lambda, d) = 0$. Suppose, on the contrary, that there exist (λ_2, d_2) and (λ'_2, d'_2) on Γ with $v(1, \lambda_2, d_2) = v(1, \lambda'_2, d'_2) = 0$. Let $J = \gamma([0, 1])$ be an arc on Γ connecting (λ_2, d_2) and (λ'_2, d'_2) with $\gamma(0) = (\lambda_2, d_2)$ and $\gamma(1) = (\lambda'_2, d'_2)$. Let γ_1 and γ_2 denote the components of γ . Because J is compact, there are only finitely many points (λ, d) in J with $v(1, \lambda, d) = 0$. Let

$$t_1 = \min\{t \in (0, 1): v(1, \gamma_1(t), \gamma_2(t)) = 0\}$$

and let $\gamma(t_1) = (\lambda''_2, d''_2)$. Then for $t \in (0, t_1)$ either $v > 0$ or $v < 0$. Let us assume, for definiteness, that for $t \in (0, t_1)$ we have $v > 0$. On $\gamma((0, t_1))$, by the implicit function theorem, we have $d = \zeta(\lambda)$, with $\zeta' > 0$. Hence $\lambda_2 \neq \lambda''_2$. Suppose $\lambda_2 < \lambda''_2$. Then taking $\lambda_n \rightarrow \lambda''_2$ with $\lambda_n < \lambda''_2$ we see that $(\lambda_n, \zeta(\lambda_n)) \rightarrow (\lambda''_2, d''_2)$ which contradicts lemma 2. Similarly, $\lambda_2 > \lambda''_2$ also leads to a contradiction. Thus, we conclude that there exists a unique $(\lambda_2, d_2) \in \Gamma$ such that $u_d(1, \lambda, d) = 0$.

Now we prove that for each $\lambda > 0$ the problem (2.1)–(2.3) has at most one stable and one unstable solution. Suppose not. Let (λ_0, d_0) and (λ_0, d'_0) be two points in Γ such that $v(1, \lambda_0, d_0) \cdot v(1, \lambda_0, d'_0) > 0$. Let $K = \psi([0, 1])$ be a path in Γ connecting (λ_0, d_0) and (λ_0, d'_0) ; we also let ψ_1, ψ_2 denote the components of ψ . Without loss of generality we can assume that ψ is one to one. Because $v(1, \lambda_0, d_0) \cdot v(1, \lambda_0, d'_0) > 0$, and ψ is one to one we see that there exists an $\varepsilon > 0$ such that $\psi_1((0, \varepsilon)) \subset (0, \lambda_0)$ or $\psi_1((0, \varepsilon)) \subset (\lambda_0, \infty)$. Let us assume that $\psi_1((0, \varepsilon)) \subset (0, \lambda_0)$; the other case can be treated in a similar way. Let t_1 be such that $\psi_1(t_1) = \min\{\psi(t): t \in (0, 1)\}$. Hence, $0 < \psi_1(t_1) < \lambda_0$. By the implicit function theorem $v(1, \psi_1(t_1), \psi_2(t_1)) = 0$. By lemma 2 there exist t_2 and t_3 near t_1 such that

$$v(1, \psi_1(t_2), \psi_2(t_2)) \cdot v(1, \psi_1(t_3), \psi_2(t_3)) < 0$$

and $t_2 < t_1 < t_3$. Without loss of generality we can assume that $v(1, \psi_1(t_2), \psi_2(t_2)) > 0$. By the intermediate value theorem there exists $t_4 \in (0, t_2)$ such that $v(1, \psi_1(t_4), \psi_2(t_4)) = 0$ which contradicts that Γ contains only one point (λ, d) with $v(1, \lambda, d) = 0$. This contradiction shows that for each λ the problem (2.1)–(2.3) has at most one stable and one unstable solution. ■

Acknowledgement—This work was partially supported by the Texas Advanced Research Program and NSF Grant DMS-8905936.

REFERENCES

1. ALI I., CASTRO A. & SHIVAJI R., Uniqueness and stability of nonnegative solutions for semipositone problems in a ball, *Proc. Am. math. Soc.* **117**(3), 775–782 (1993).
2. CASTRO A. & SHIVAJI R., Nonnegative solutions for a class of nonpositone problems, *Proc. R. Soc. Edinb.* **108A**, 291–302 (1988).
3. CASTRO A., GARNER J. & SHIVAJI R., Existence results for classes of sublinear semipositone problems, *Result. Math.* **23**, 214–220 (1993).
4. CASTRO A. & SHIVAJI R., Nonnegative solutions for a class of radially symmetric nonpositone problems, *Proc. Am. math. Soc.* **106**(3), 735–740 (1989).
5. GIDAS B., NI W. & NIRENBERG L., Symmetry and related properties via the maximum principles, *Communs math. Phys.* **68**, 209–243 (1979).
6. PROTTER M. H. & WEINBERGER H. F., *Maximum Principles in Differential Equations*. Prentice-Hall, Englewood Cliffs, New Jersey (1967).